

## RELATIONSHIP OF THE PHYSICO-CHEMICAL CONSTANTS WITH PERIODIC LAW

Vassiliev V.P., Lysenko V.A.

*Lomonosov Moscow State University, Chemistry Department, 119991 Moscow, Russia  
e-mail: valeryvassiliev@yahoo.fr*

Different relationships between the various physical and chemical properties of isostructural compounds take place according to the Periodic Law that is a fundamental basis of Chemistry. Previous review articles evidenced the relation of thermodynamic data with Periodic Law [1-4] and established a strict relationship between the enthalpy of formation, melting point and the atomic numbers of components in the semiconductor A<sup>III</sup>B<sup>V</sup> phases, with diamond-like structure of sphalerite and wurtzite types. The proposed model was used for the critical assessment of the thermodynamic properties of isostructural compounds. The relationship between the reduced enthalpy ( $\Delta_f H_{298}^{\circ}/T_m$ ), standard entropy ( $S_{298}^{\circ}$ ), reduced Gibbs energy and the sum of the atomic numbers ( $Z_i = Z_A + Z_B$ ) has been used for a critical assessment of the thermodynamic properties of A<sup>III</sup>B<sup>V</sup> phases.

The Similarity Method was used for the critical analysis of specific heats  $C_p^{\circ}(T)$  for solid state of the A<sup>III</sup>B<sup>V</sup> isostructural phases. A critical analysis of heat capacities was carried out for the pure elements of the Periodic System fourth group (C, Si, Ge, Sn) and isostructural phases A<sup>III</sup>B<sup>V</sup> and A<sup>II</sup>B<sup>VI</sup>. It was found that the dependence of the heat capacities from 0 to 1500 K follows certain regularity. Phases with the same sum of the atomic numbers of elements ( $Z_i$ ), such as BN (hex)  $Z_i = 12$  and glassy pure carbon  $Z = 6$ ; BP and AlN ( $Z_i = 20$ ); AlP ( $Z_i = 28$ ) and pure Si ( $Z = 14$ ); BAs and GaN ( $Z_i = 38$ ); AlAs and ZnS ( $Z_i = 46$ ); AlSb, GaAs, InP, CDs ( $Z_i = 64$ ) and pure Ge ( $Z = 32$ ); GaSb, InAs, and CdSe ( $Z_i = 82$ ); InSb, CdTe ( $Z_i = 100$ ) and pure grey Sn ( $Z = 50$ ); have the same heat capacity experimental values of the solid state within the experimental uncertainty.

The second group is concerned of the rare earth metals and their compounds and the actinides. The "tetrad-effect" phenomenon was established and used for the analysis, correction and prediction of thermodynamic data for the lanthanide (Ln) compounds and the pure actinides (Ac). They are connected to the 4f-electrons of the lanthanide elements (Ln: La-Lu; their atomic numbers are 57-71) and 5f-electrons of the actinides with atomic numbers from 89 to 102. The most sensitive to the tetrad-effect the thermodynamic functions of lanthanide and actinide compounds are standard entropies and entropies of formation, because they are the most susceptible to the influence of the 4f- and 5f electrons of the lanthanides and the actinides. We analyzed some classes of lanthanides with other elements of the Periodic Table, which account for only a sampling of the thousands of similar binary compounds possible. As an example, we use the tetrad-effect concept for the analysis and prediction of the standard entropies of the solid phases as the Ln<sub>2</sub>X<sub>3</sub> (X=O, S, Se, Te) [5]. This approach can also be applicable to other classes of the Ln compounds as LnN, LnB<sub>2</sub>, LnB<sub>4</sub>, LnB<sub>6</sub>, LnF<sub>3</sub>, LnIn<sub>3</sub> [6, 7] and other compounds. Unfortunately, the actinides and the alloys on the base these elements are studied insufficiently, but it is possible to apply to them the same laws.

**Keywords:** Semiconductor (AII-BVI, AIV, AIII-BV) and lanthanide compounds, thermodynamic properties, entropy, Debye's functions, crystal structure, critical assessment, tetrad-effect

### Literature

- [1] V.P. Vassiliev, B. Legendre, V.P. Zlomanov, The critical analysis and mutual coherence of thermodynamic data of the A<sup>III</sup>B<sup>V</sup> phases, *Intermetallics* 19, 1891 (2011).
- [2] Vassiliev V.P., Taldrik A.F., Ilinykh N.I., New Correlative Method of Thermodynamic Analysis of the Inorganic Compounds. MATEC Web of Conferences, № 01078 (2013)
- [3] V.P. Vassiliev, W. Gong, A.F. Taldrik, S.A. Kulinich, Method of the correlative optimization of heat capacities of isostructural compounds, *J. Alloys Comp.* 552, 248 (2013)
- [4] V.P. Vassiliev, Lysenko V.A., Taldrik A.F., Ilinykh N.I., Sevastyanova L.G. Approximation of the low-temperature heat capacity of A<sup>III</sup>B<sup>V</sup> compounds by linear combination of Debye's functions. 41st Conference on Phase Equilibria Book of Abstracts, JEEP, Coimbra, Portugal, March 23-25 140 (2015)
- [5] V.P. Vassiliev, and V.A. Lysenko, A new approach for the study of thermodynamic properties of lanthanide compounds, *Electrochim. Acta* 222, 1770 (2016).
- [6] V.P. Vassiliev, A.F. Taldrik., B. Legendre, Analyse thermodynamique des terres rares et leurs alliages REIn<sub>3</sub>, *JEEP* 2009, DOI: 10.1051/jeep/200900004
- [7] V.P. Vassiliev, B. Ablazeze, and A.F. Taldrik, Thermodynamic analysis of the rare earth metals and their alloys with indium in solid state, *J. Alloys Comp.* 572, 118 (2013).